

Alcohol Consumption Demographics

Table 9. Percent (standard error) of women who reported drinking alcohol during the 3 months before pregnancy. Ques. # 32a

Maternal Characteristics	percent (s.e.)
Total	54.4(1.4)
Race	
Black	40.4(2.4)
White	63.4(1.7)
Other	39.6 (9.6)
Age	
Less than 20	36.5(3.9)
20-29	57.0(1.9)
30+	57.0(2.6)
Education	
Less than High School	38.2(3.4)
High School	52.1(2.4)
More than High School	63.3(2.0)
Marital Status	
Married	56.6(1.8)
Other	51.7(2.3)
Birthweight	
LBW (<2500 grams)	48.4(4.1)
NBW (>= 2500 grams)	55.1(1.5)
Medicaid Status*	
Medicaid	50.2(1.9)
Other	61.2(2.1)

Table 10. Percent (standard error) of women who reported drinking alcohol during the last trimester of pregnancy, according to sociodemographic characteristics. Ques. # 33a

Maternal Characteristics	percent (s.e.)
Total	6.8(0.7)
Race	
Black	3.5(0.9)
White	9.1(1.0)
Other	0.0
Age	
Less than 20	6.4(2.1)
20-29	4.9(0.8)
30+	10.8(1.5)
Education	
Less than High School	5.4(1.5)
High School	6.1(1.1)
More than High School	8.0(1.1)
Marital Status	
Married	7.8(1.0)
Other	5.6(1.0)
Birthweight	
LBW (<2500 grams)	6.2(1.8)
NBW (>= 2500 grams)	6.9(0.8)
Medicaid Status*	
Medicaid	8.9(1.2)
Other	5.5(0.8)

* A mother is considered "on Medicaid" if she was enrolled in Medicaid during any of the following time periods: just before pregnancy, during prenatal care or during delivery.